

Problem 1. (24 points)

Clearly explain your reasoning. **There are questions on the back of this page.**

- (a) [6 points] Explain why we say there is no such thing as a classical free particle in quantum mechanics. How do we describe a “free particle” in quantum mechanics?

- (b) [6 points] For the half harmonic oscillator potential

$$V(x) = \begin{cases} \infty, & x \leq 0 \\ \frac{1}{2}m\omega^2x^2, & x > 0 \end{cases}$$

what are the energies of the two lowest states? Express your answer in terms of m , ω , and constants, as appropriate.

Problem 1 continued:

- (c) [4 points] Let $\{\psi_n\}$ be the set of wavefunctions that are solutions to the harmonic oscillator potential with frequency ω for a particle of mass m . Consider the wavefunction

$$\psi(x) = A(3\psi_1 + 4\psi_3).$$

Find A .


- (d) [4 points] For the wave function in the previous part what is $\langle \hat{x} \rangle$? [*Hint:* The answer can be calculated without doing any integrals. One way to do this is to use the symmetries of the wave functions.]

- (e) [4 points] For this wave function what is the expectation value of the energy, $\langle \hat{H} \rangle$? Express your answer in terms of m , ω , and constants, as appropriate.

Problem 2. (20 points)

A particle of mass m is in an infinite square well given by the potential

$$V(x) = \begin{cases} \infty, & x \leq 0 \\ 0, & 0 < x < a \\ \infty, & x \geq a \end{cases} .$$



The particle is placed in the left half of the well with the initial state

$$\Psi(x, 0) = \begin{cases} 0, & x \leq 0 \\ A, & 0 < x < a/2 \\ 0, & x \geq a/2 \end{cases} .$$

(a) [4 points] Calculate the value of A .

(b) [10 points] Expand the initial state in terms of the solutions to the time independent Schrödinger equation, $\psi_n(x)$. [Note: Determining the expansion coefficients is sufficient.]

(c) [6 points] Write down the energy of the ground state, and first, second, and third excited states. Also write down the probability of finding the particle in each of these states.


Problem 3. (26 points)

Consider the potential given by

$$V(x) = -\alpha\delta(x) + V_0\theta(x)$$

where $\theta(x)$ is the step function (it is 1 when $x \geq 0$, zero otherwise), as shown at the right.

[Note: **There are questions on the back of this page.**]



- (a) [2 points] What condition must be satisfied by the energy for this potential to have bound states?
- (b) [5 points] Write down the bound state wavefunctions in the region $x < 0$ and $x > 0$. Clearly define any constants you introduce.
- (c) [5 points] Use the continuity of the wave function to solve for the wave function up to an overall normalization factor.
- (d) [5 points] Without doing a calculation do we expect $\langle \hat{x} \rangle < 0$, $\langle \hat{x} \rangle = 0$, or $\langle \hat{x} \rangle > 0$ for the bound state? Justify your answer.

Problem 3 continued:

(e) [6 points] Find an equation that determines the energy of the bound state in terms of m , α , E , V_0 , and constants, as necessary. Do **not** solve the equation, leave it in a clear form.

(f) [3 points] How many bound states are there? Solve for the energy of the bound states.